

Lectotypification of Brazilian species of *Vitex* Tour. ex L. (Lamiaceae) Lectotipificação de espécies brasileiras de *Vitex* Tour. ex L. (Lamiaceae)

Flávio França¹

Abstract: Lectotypes are designated for the following *taxa* of *Vitex* Tour. ex L. (Lamiaceae) from Brazil: *Vitex brevilabiata* Ducke, *V. duckei* Huber, *V. gardneriana* Schauer, *V. schaueriana* Moldenke, *V. triflora* Vahl var. *angustiloba* Huber, *V. triflora* var. *kraatzii* Huber and *V. triflora* var. *tenuiflora* Huber.

Keywords: *Vitex*. Lamiaceae. Verbenaceae. Brazil.

Resumo: Lectótipos são designados para os seguintes *taxa* de *Vitex* Tour. ex L. (Lamiaceae) do Brasil: *Vitex brevilabiata* Ducke, *V. duckei* Huber, *V. gardneriana* Schauer, *V. schaueriana* Moldenke, *V. triflora* Vahl var. *angustiloba* Huber, *V. triflora* var. *kraatzii* Huber e *V. triflora* var. *tenuiflora* Huber.

Palavras-chave: *Vitex*. Lamiaceae. Verbenaceae. Brasil.

¹ Universidade Estadual de Feira de Santana. Departamento de Ciências Biológicas. Feira de Santana, Bahia, Brasil (flaviofranca@hotmail.com).



INTRODUCTION

The name *Vitex* was first used by Tournefort (1719), when he described five species. Linnaeus (1753) included four species in the genus. Almost a hundred years later, Schauer (1847) published a global revision of *Vitex*, in which he listed 59 species. In his contribution to "Flora Brasiliensis", Schauer (1851) included ten species of *Vitex*, later, Moldenke (1959) recognized 36 species for Brazil.

An inventory of Brazilian *Vitex* species (in preparation) has revealed that some names have no designed holotype, and this paper aims to lectotypify these.

RESULTS

Vitex breviliabiata Ducke, Arch. Jard. Bot. Rio de Janeiro 4: 172-173. 1925. Brazil, Pará, Tapajós river, Poçoão, "habitat in silvis non inundatis regione cataractarum inferiorum fluminis Tapajóz", 07.12.1919 (fl) A. Ducke s.n. RB 311 (RB, Lectotype, here designated).

This species name is based on two sheets (A. Ducke s.n. RB 18954 and A. Ducke s.n. RB 311) both deposited in the herbarium of the Rio de Janeiro Botanical Garden – RB – (Ducke, 1925). There are duplicates of A. Ducke s.n. RB 18954 at Kew (K). The sheet RB 311 was selected as lectotype because its leaves are more developed than the sheet 18954.

Vitex duckei Huber, Bol. Mus. Para. Emílio Goeldi 5: 217-218. 1908. Brazil, Pará, "Habitat in locibus campestribus ad flumen Yamundá (Nhamundá) inferius prope Faro", 04.09.1907 (fl), A. Ducke s.n. MG 8605 (MG, Lectotype here designated).

Huber (1908) described *V. duckei* based on two specimens: MG 8441 (MG, RB) and MG 8605 (MG). MG 8441 sheet has few flowers, but MG 8605 contains many flowers, with buds and opened flowers. This sheet is therefore designated here as the lectotype for this species.

Vitex gardneriana Schauer, in De Candolle. Prodr., 11: 687-688, 1847. Brazil, Pernambuco, "in rivis Capiberibe, Moxoto", 1842 (fl), Houillet s.n. (BR, Lectotype here designated).

Schauer (1847) described *V. gardneriana* based on two collections, one of Gardner and another of Houillet. He mistakenly said that the specimens were collected in Rio de Janeiro, but he stated, in "Flora Brasiliensis" (Schauer, 1851), that the material was collected at "rivis Capiberibe, Moxoto", a river located in Pernambuco.


The material deposited at the herbarium at Kew (K) under Gardner's number (Gardner 1107, two sheets) was designated as 'cotype' of *V. gardneriana* by Moldenke. However, the protologue lists only a numberless Gardner collection, so the material from Kew is apparently not the type of the species.

Schauer (1847) indicated that he saw the material in the herbaria of Martius and De Candolle. The specimen from Geneva (G), where the De Candolle collections are deposited was not examined. Only the Houillet material deposited at BR, where are the Martius collections, was studied here.

The Houillet specimen from BR display clearly the principal features of the species and includes the citation of Schauer in "Flora Brasiliensis", "in rivis Capiberibe Moxoto". Based on this, it is designated here as the lectotype of this species.

Vitex schaueriana Moldenke, Revista Sudamer. Bot. 5: 3. 1937. ≡ *V. montevidensis* var. *parviflora* Schauer, in De Candolle. Prodr 11: 689. 1847. Brazil, Bahia, Serra de Açuruá, 1839 (fl), J. S. Blanchet 2782 (W, Lectotype, here designated. Isolectotype: K).

Schauer (1847) described *V. montevidensis* var. *parviflora* based on two collections: J. S. Blanchet 2782 and Riedel s.n. The Riedel's specimen could not be found, but the one of J. S. Blanchet 2782 is well represented. There are duplicates of J. S. Blanchet 2782 in Kew (K) and Vienna (W). Schauer (1847) cited collections deposited in Berlin (B), Vienna (W) and Saint Petersburg (LE) ("berol. c. vindob. et petr."). The only material cited by Schauer (1847) examined was the one from Vienna (W), however, it matches the original description of this species. Additionally, a label giving the correct locality data is given. Therefore this specimen is designated here as the lectotype of *Vitex montevidensis* var.



parviflora and *V. schaueriana*, name proposed by Moldenke (1937), because there is an earlier species named *V. parviflora*.

Vitex triflora Vahl var. *angustiloba* Huber, Bol. Mus. Para. Emílio Goeldi, 5(1): 215. 1908. Brazil, Pará, Soure, São Salvador, 30.11.1904 (fl), R. Siqueira s.n. MG 6886 (MG, Lectotype, here designated).

Huber (1908) described this variety based on three specimens: Huber s.n. MG 254 (MG, NY – fragments-), Huber s.n. MG 524 (MG, RB) and R. Siqueira s.n. MG 6886. The latter specimen (R. Siqueira s.n. MG 6886) is designated as lectotype, because it is the most representative, clearly showing the variety characters mentioned in the protologue.

Vitex triflora Vahl var. *kraatzii* Huber, Bol. Mus. Para. Emílio Goeldi 5: 216. 1908. Brazil, Pará: Monte Alegre, field, jul. 1902 (fl), A. Ducke s.n. MG 2873 (MG, Lectotype, here designated).

This variety was described by Huber (1908) based on two sheets: Huber s.n. MG 1729 and A. Ducke s.n. MG 2873. The former has some damage to its leaves, the latter is better conserved. Therefore, the specimen A. Ducke s.n. MG 2873 is designated here as the lectotype of this variety.

Vitex triflora Vahl var. *tenuifolia* Huber, Bol. Mus. Para. Emílio Goeldi 5: 214. 1908. Brazil, Amazonas: Rio Japurá, “logar jupará”, capueira, sept. 1904 (fl), A. Ducke s.n. MG 6759 (MG, Lectotype, here designated).

Huber (1908) described this variety based on nine specimens, all collected by A. Ducke: MG 1890, MG 2872, MG 3758, MG 4927, MG 2908a, MG 7914, MG 8381, MG 6759, and MG 2950b. The specimen A. Ducke s.n. MG 6759 has the main characters for the recognition of the variety and, therefore, it is designated here as lectotype of *V. triflora* var. *tenuifolia*.

ACKNOWLEDGMENTS

This work was partially supported by Mellon Foundation/Margaret Mee and MCT/PPBio-Semi-árido Grant. I thanks Raymond Harley and Cássio van den Berg for helping with the English language and nomenclatural aspects.

REFERENCES

- DUCKE, A., 1925. Plantes nouvelles ou peu connues de la région amazonienne III. **Archivos do Jardim Botânico do Rio de Janeiro** 4: 172-174.
- HUBER, J., 1908. As espécies amazônicas do gênero *Vitex*. **Boletim do Museu Paraense Emílio Goeldi** 5: 209-222.
- LINNAEUS, C., 1753. *Vitex*. In: C. LINNAEUS (Ed.): **Species Plantarum**: 638 (938). Laurentii Salvii, Holmiae.
- MOLDENKE, H. N., 1937. Some needful nomenclatural changes. **Revista Sudamericana de Botanica** 5: 1-3.
- MOLDENKE, H. N., 1959. **A Résumé of the Verbenaceae, Avicenniaceae, Stilbaceae, Symphoremaceae and Eriocaulaceae of the world as to valid taxa, geographic distribution and synonymy**: 1-495. H. N. Moldenke, New Jersey.
- SCHAUER, J. C., 1847. Verbenaceae. In: A. P. DE CANDOLE (Ed.): **Prodromus Systematis Naturalis Regni Vegetabilis** 11: 522-701.
- SCHAUER, J. C., 1851. Verbenaceae. In: C. F. P. VON MARTIUS & A. G. EICHLER (Eds.): **Flora Brasiliensis** 9: 169-308. Typographia Regia, Munchen, Wien, Leipzig.
- TOURNEFORT, J. P., 1719. **Institutiones Rei Herbariae**: 1: 603. Editio Tertia, Paris.

Recebido: 14/07/2009

Aprovado: 21/11/2010

Responsabilidade editorial: Anna Luiza Ilkiu Borges

