










Intestinal parasitism in traditional communities surrounding an environmental protection area in Maranhão State, northeastern Brazil

Parasitismo intestinal em comunidades tradicionais vivendo ao redor de uma área de proteção ambiental no estado do Maranhão, nordeste do Brasil

Maria Lindalva Alves da Silva¹, Célia Mariza de Oliveira Guerra Álvares¹, Jurecir da Silva², Julio Cesar Pegado Bordignon¹, Mayron Moraes Almeida¹, Eduilson Lívio Neves da Costa Carneiro², Filipe Anibal Carvalho-Costa³, Kerla Joeline Lima Monteiro⁴, Antonio Henrique Almeida de Moraes Neto¹

¹ Fundação Oswaldo Cruz, Instituto Oswaldo Cruz, Laboratório de Inovações em Terapias, Ensino e Bioprodutos, Rio de Janeiro, Rio de Janeiro, Brasil

² Instituto Federal de Educação, Ciência e Tecnologia do Piauí, Teresina, Piauí, Brasil

³ Fundação Oswaldo Cruz, Instituto Oswaldo Cruz, Laboratório de Epidemiologia e Sistemática Molecular, Rio de Janeiro, Rio de Janeiro, Brasil

⁴ Fundação Oswaldo Cruz, Escritório Regional do Piauí, Teresina, Piauí, Brasil

ABSTRACT

OBJECTIVES: To estimate prevalence and describe socio-ecological factors associated with intestinal parasitism in rural and peri-urban communities surrounding an environmental protection area in Maranhão State, Brazil. **MATERIALS AND METHODS:** A cross-sectional survey was conducted on 469 subjects in two communities around the Chapada das Mesas National Park. Qualitative and quantitative parasitological examinations were performed, and socio-ecological data were assessed. A geographic information system was used for spatial analysis. **RESULTS:** The prevalence of hookworm infection was 20.6% in Canto Grande and 2.4% in Alto Bonito. The ascariasis prevalence in Canto Grande and Alto Bonito was 0.4% and 2.9%, respectively. Infections with gut protozoa were more frequent in Alto Bonito. Hookworm infection was associated with the male sex, open defecation, clay floors in homes, and lower monthly family income. Ascariasis was associated with open defecation. Ingestion of water directly from a river was associated with *Giardia duodenalis* infection. A total of 86.0% of the hookworm-positive subjects had light infection intensity, 4.0% moderate, and 10.0% heavy. Geoprocessing demonstrated that soil-transmitted helminths hotspots were in Canto Grande, while protozoa were concentrated in Alto Bonito. **CONCLUSION:** The present study suggests that intestinal parasitism should be considered a heterogeneous group of infections with distinct determinants in the studied communities. Intestinal parasitism persists, revealing the inadequacy of sanitation in traditional communities around the Chapada das Mesas National Park.

Keywords: Intestinal Parasitism; Chapada das Mesas National Park; Soil-Transmitted Helminthiasis.

RESUMO

OBJETIVOS: Estimar a prevalência e descrever os fatores socioecológicos associados ao parasitismo intestinal em comunidades rurais e periurbanas ao redor de uma área de proteção ambiental no estado do Maranhão, Brasil. **MATERIAIS E MÉTODOS:** Estudo transversal, realizado com 469 indivíduos de duas comunidades do entorno do Parque Nacional da Chapada das Mesas. Exames parasitológicos qualitativos e quantitativos foram realizados, e dados socioecológicos foram avaliados. Um sistema de informações geográficas foi utilizado para a análise espacial. **RESULTADOS:** A prevalência de ancilostomíase foi de 20,6% em Canto Grande e 2,4% em Alto Bonito. A prevalência de ascaridíase em Canto Grande e Alto Bonito foi de 0,4% e 2,9%, respectivamente. Infecções por protozoários intestinais foram mais frequentes em Alto Bonito. A ancilostomíase foi associada ao sexo masculino, defecação a céu aberto, pisos de barro nas casas e menor renda familiar mensal. A ascaridíase foi associada à defecação a céu aberto. A ingestão de água diretamente de um rio foi associada à infecção por *Giardia duodenalis*. Um total de 86,0% dos indivíduos positivos para ancilostomídeos apresentaram intensidade de infecção leve, 4,0% moderada e 10,0% intensa. O geoprocessamento demonstrou que os hotspots de helmintos transmitidos pelo solo estavam em Canto Grande, enquanto os protozoários estavam concentrados em Alto Bonito. **CONCLUSÃO:** O parasitismo intestinal deve ser considerado um grupo heterogêneo de infecções com determinantes distintos nas localidades estudadas. Sua persistência revela a inadequação do saneamento nas comunidades tradicionais do entorno do Parque Nacional da Chapada das Mesas.

Palavras-chave: Parasitismo Intestinal; Parque Nacional da Chapada das Mesas; Helmintíases Transmitidas pelo Solo.

Correspondence / Correspondência:

Antonio Henrique Almeida de Moraes Neto

Fundação Oswaldo Cruz, Instituto Oswaldo Cruz, Laboratório de Inovações em Terapias, Ensino e Bioprodutos

Av. Brasil, 4365. Bairro: Manguinhos. Zip code: 21040-900 – Rio de Janeiro, Rio de Janeiro, Brasil – Phone #: +55 (21) 2598-4220

E-mail: ahmn@ioc.fiocruz.br

INTRODUCTION

In the Brazilian backlands, many rural communities maintain their traditional lifestyle. The Chapada das Mesas National Park is an environmental protection area in Maranhão State, within the Cerrado biome¹, and families residing around the Park maintain their cultural traditions as defined by the National Policy for Sustainable Development of Traditional Peoples and Communities^{2,3}. The communities surrounding the Park are characterized by the sustainable use of natural resources and by maintaining their cultural traditions through their customs and beliefs⁴. These populations practice subsistence agriculture, characterized by family gardens of beans, corn, cassava, and bananas, and palm fruit extraction, such as buriti (*Mauritia flexuosa*) and bacuri (*Platonia insignis*)⁵. They also raise animals, such as chickens and pigs. Nevertheless, this population has very few sources of income, and 75% of the families survive with less than the Brazilian minimum wage, with a large proportion of them benefiting from governmental income transfer programs, which characterizes an economic poverty scenario⁵. The sanitation systems are very incipient⁶, and waste management has been a challenge, which favors the spread of parasitic diseases. For many reasons, open defecation is still practiced⁶, leading to significant environmental contamination with the infective stages of intestinal parasites. The transmission of waterborne diseases in these communities is also favored by the consumption of water collected directly from rivers, not subjected to any treatment or purification system⁶.

Intestinal parasitism encompasses a set of infections caused by distinct organisms, which can be helminths or protozoa⁷. Among gut parasites, soil-transmitted helminths (STH) spend a phase of their life cycle in the soil, which is mandatory for the maturation of the infective stages. These parasites exist in the soil as rhabditoid and filariform larvae for hookworms (*Necator americanus* and *Ancylostoma duodenale*) and embryonated eggs for *Ascaris lumbricoides* and *Trichuris trichiura*⁸. These organisms, therefore, have different transmission routes, being percutaneous for hookworms and oral for *A. lumbricoides* and *T. trichiura*, leading to different dissemination dynamics. Parasitic gut protozoa are transmitted orally through the ingestion of infective cysts. Among them, *Giardia duodenalis* and *Entamoeba histolytica* are the most important due to their pathogenic potential, including malabsorption, diarrhea, and dysentery^{9,10}. Some protozoa are considered human gut commensals, such as *Entamoeba coli*, *Endolimax nana*, and *Iodamoeba bütschlii*¹¹. Percutaneous transmission organisms may have a more spatially restricted cycle in the peri-domestic environment, associated with soil contamination caused by local open defecation and poor sanitation¹². Orally transmitted parasites may be disseminated over longer distances by contaminated water and food due to the resistance and longevity of their eggs and cysts.

Control strategies have shaped the etiological profile of intestinal parasitism in recent decades. Mass drug administration (MDA) with anthelmintics has reduced

the prevalence and parasitic load of STH, leading to an increase in the participation of protozoa in the species composition. This shift occurs because anthelmintic drugs, such as 400 mg single-dose albendazole, do not target protozoa^{13,14}. Additionally, standard Kato-Katz smears routinely utilized in population surveys are unable to detect the cysts of gut parasitic protozoa, further exacerbating the neglect of these infections.

Maranhão State, in northeastern Brazil, has one of the highest prevalence of STH in the country, with 15.8% for hookworm, 17.5% for *A. lumbricoides*, and 5.8% for *T. trichiura* in children aged 7 to 17 years¹⁵. Hookworm infection and ascariasis presented prevalence rates of 14.3% and 9.3%, respectively, in studies conducted with school-aged children living in peri-urban communities in Maranhão¹⁶. These data suggest that rural and peri-urban communities in the State are particularly vulnerable to intestinal parasitism. Previous studies carried out in communities around the Chapada das Mesas National Park^{4,5,6} motivated the investigation of the epidemiological aspects of intestinal parasitism in the region to describe the socio-ecological determinants of this group of infections. The etiological profile of gut parasitic infections and the parasite load of STH are important for planning improvements in control strategies. The present study aimed to estimate prevalence and describe socio-ecological factors associated with intestinal parasitism in rural and peri-urban communities surrounding an environmental protection area in Maranhão State, Brazil.

MATERIALS AND METHODS

SETTING

The study was carried out in two traditional communities living around the Chapada das Mesas National Park, established by law in 2005¹: the community of Canto Grande (population of 1,544 inhabitants; located at 7°06'30"S/48°13'33"W), in Carolina City; and the community of Alto Bonito (population of 2,148 inhabitants; located at 7°23'30"S/46°37'21"W), in Riachão City. Canto Grande is predominantly composed of rural communities, and Alto Bonito, of peri-urban neighborhoods (Figure 1).

STUDY DESIGN, SAMPLING STRATEGY, AND STATISTICAL ANALYSES

A cross-sectional survey was conducted in July 2022. Among the 335 subjects visited in Canto Grande, 262 (78.2%) returned a fecal sample. In Alto Bonito, from 287 people invited to participate in the study, 207 (72.1%) returned a fecal sample. In both communities, socio-ecological and demographic data were obtained through interviews carried out by researchers during home visits. These data included income, sanitation and housing characteristics, practice of open defecation, and source of drinking water. Residents were selected with the support of Community Health Agents (CHA). The sample size followed the parameters: margin of error of 5%, confidence level of 95%, and expected frequency of 20% for infection with gut parasites.




Figure 1 – Map of the studied area

The results were presented as frequencies of infection by different parasite species, which were compared using prevalence ratios (PR) and respective 95% confidence intervals (CI). Statistical significance was assessed using Fisher's test. The houses visited were georeferenced using GPS, and QGIS¹⁷ was used for spatial analysis. The base map was acquired from IBGE (Brazilian Institute of Geography and Statistics), and the coordinates of the sampling sites were recorded in the SAD 69 datum geodetic coordinate system. For exploration and modeling, the maps were analyzed for spatial point patterns using the kernel method. A 2,000 m search radius was selected, and raster output maps were generated with a pixel size of 10 x 10 m. Density values on the kernel density map were scaled to visually represent individual kernels.

PARASITOLOGICAL ANALYSES

The Lutz technique was used to detect helminth eggs and protozoan cysts. For STH quantitative analysis, the Kato-Katz technique was used (Helm Test[®] kit, LabHouse, Belo Horizonte, Minas Gerais, Brazil). Infection intensity was determined as follows: (i) hookworm infection – light, with a parasitic load of $\leq 1,999$ eggs per gram of feces (epg); moderate, with a load between 2,000 and 3,999 epg; and heavy, with $\geq 4,000$ epg; and (ii) ascariasis – light, ranged of 0 to 4,999 epg; moderate, from 5,000 to 49,999 epg; and heavy, with $\geq 50,000$ epg¹⁸.

ETHICS

This study received approval from the Ethics Committee for Research with Humans at the Oswaldo

Cruz Institute, Oswaldo Cruz Foundation (IOC/Fiocruz), under CAAE No. 40483120.7.0000.5248 (23 December 2020), and authorization from the Chico Mendes Institute for Biodiversity Conservation (SISBIO/ICMBio), under No. 79178-1 (9 June 2022).

RESULTS

Among the 469 residents studied in the two communities, 42.6% were classified as living in poverty (family income less than the minimum wage), 31.6% practiced open defecation, 16.2% lived in houses with uncoated clay floors, and 4.1% drank water directly from the river (Table 1).

Differences were identified in the etiological profile of intestinal parasitism in the two communities. The hookworm infection prevalence was 20.6% in Canto Grande and 2.4% in Alto Bonito (PR=8.56; 95% CI=3.49–21.02; $p < 0.001$). The ascariasis prevalence in Canto Grande and Alto Bonito was 0.4% and 2.9%, respectively (PR=0.13; CI 95%=0.02–1.08; $p=0.047$). Infections with gut protozoa were more frequent in Alto Bonito, and statistically significant differences were observed for *Entamoeba coli* and *E. nana*. The factors associated with STH infections are also shown in table 1. Specifically, hookworm infection was associated with the male sex, open defecation, clay floors in homes, and lower monthly family income. Ascariasis was also associated with open defecation. Concerning intestinal protozoa, the ingestion of water directly from a river was associated with *G. duodenalis* infection (Table 2).

Table 1 – Prevalence of soil-transmitted helminthiases in different groups defined by sociodemographic variables in Canto Grande and Alto Bonito communities, Chapada das Mesas National Park, Maranhão State, Brazil, 2022


Variables	Total samples		Hookworm infection			Ascariasis				
	N	%	N	%	PR (95% CI)	p-value	N	%	PR (95% CI)	p-value
Community										
Alto Bonito	207	44.1	5	2.4	1		6	2.9	1	
Canto Grande	262	55.9	54	20.6	8.56 (3.49–21.02)	< 0.001	1	0.4	0.13 (0.02–1.08)	0.047
Age group										
< 10 years	85	18.1	9	10.6	1		1	1.2	1	
10 – 17 years	54	11.5	10	18.5	1.72 (0.75–3.97)	0.213	4	7.4	6.29 (0.72–54.84)	0.074
≥ 18 years	330	70.4	40	12.1	1.13 (0.57–2.23)	0.850	2	0.6	0.51 (0.04–5.61)	0.498
Sex										
Female	243	51.8	22	9.1	1		–	–	1	
Male	226	48.2	37	16.4	1.80 (1.09–2.95)	0.025	7	3.1	Undefined	0.005
Income (R\$)										
> 1,300 BRL	269	57.4	27	10.0	1		3	1.1	1	
≤ 1,300 BRL	200	42.6	32	16.0	1.75 (1.08–2.83)	0.023	4	2.0	1.80 (0.40–7.99)	0.464
Defecation site										
Latrine	321	68.4	13	4.0	1		1	0.3	1	
Open (bushes)	148	31.6	46	31.1	7.65 (4.26–13.71)	< 0.001	6	4.1	13.01 (1.58–107.13)	0.004
House floor										
Coated	393	83.8	29	7.4	1		2	0.5	1	
Clay	76	16.2	30	39.5	5.33 (3.41–8.34)	< 0.001	5	6.6	12.92 (2.55–65.41)	< 0.001
Drinking water source										
Piped	397	84.6	57	14.4	1		7	1.8	1	
Rudimentary well	36	7.7	–	–	Undefined	0.008	–	–	Undefined	1.000
Water truck	17	3.6	–	–	Undefined	0.143	–	–	Undefined	1.000
Direct from river	19	4.1	2	10.5	0.73 (0.19–2.77)	1.000	–	–	Undefined	1.000

Conventional sign used: – Data equal to zero, not resulting from rounding; PR: Prevalence ratio; 95% CI: Confidence intervals.

Table 2 – Prevalence of intestinal protozoan infections in different groups defined by sociodemographic variables in Canto Grande and Alto Bonito communities, Chapada das Mesas National Park, Maranhão State, Brazil, 2022

Variables	Total samples			<i>Giardia duodenalis</i>			<i>Entamoeba histolytica/dispar</i>			<i>Entamoeba coli</i>			<i>Entodolimax nana</i>				
	N	%	N	%	PR (95% CI)	p-value	N	%	PR (95% CI)	p-value	N	%	PR (95% CI)	p-value			
Community																	
Alto Bonito	207	44.1	19	9.2	1	0.096	7	3.4	1	0.156	30	11.5	1	60	29.0	1	
Canto Grande	262	55.9	13	5.0	0.54 (0.27–1.07)	0.096	4	1.5	0.45 (0.13–1.52)	0.156	49	23.7	0.48 (0.32–0.73)	28	10.7	0.37 (0.24–0.55)	< 0.001
Age group																	
< 10 years	85	18.1	9	10.6	1	1.000	2	2.4	1	1.000	18	21.2	1	16	18.8	1	
10–17 years	54	11.5	5	9.3	0.87 (0.30–2.47)	1.000	–	–	Undefined	0.512	8	14.8	0.69 (0.32–1.49)	14	25.9	1.38 (0.73–7.58)	0.398
≥ 18 years	330	70.4	18	5.5	0.51 (0.24–1.11)	0.134	9	2.7	1.16 (0.25–5.26)	1.000	53	16.1	0.75 (0.47–1.22)	58	17.6	0.93 (0.56–1.54)	0.753
Sex																	
Female	243	51.8	14	5.8	1	0.365	5	2.1	1	0.765	44	18.1	1	51	21.0	1	
Male	226	48.2	18	8.0	1.38 (0.70–2.71)	0.365	6	2.7	1.29 (0.39–4.16)	0.765	35	15.5	0.85 (0.57–1.28)	37	16.4	0.78 (0.53–1.14)	0.236
Income (R\$)																	
≤ 1,300 BRL	200	42.6	15	7.5	1	0.711	2	1	1	0.126	33	16.6	1	38	19.1	1	
> 1,300 BRL	269	57.4	17	6.3	0.84 (0.43–1.64)	0.711	9	3.3	3.34 (0.73–15.31)	0.126	46	17.0	1.03 (0.68–1.55)	50	18.5	0.97 (0.66–1.43)	0.905
Defecation site																	
Latrine	321	68.4	21	6.5	1	0.698	11	3.4	1	0.020	54	16.8	1	61	19.0	1	
Open (bushes)	148	31.6	11	7.4	1.13 (0.56–2.29)	0.698	–	–	Undefined	0.020	25	16.9	1.00 (0.92–1.09)	27	18.2	0.99 (0.90–1.09)	0.476
House floor																	
Coated	393	83.8	24	6.1	1	0.209	11	2.8	1	0.139	63	16.0	1	71	18.1	1	
Clay	76	16.2	8	10.5	1.72 (0.80–3.69)	0.209	–	–	Undefined	0.139	16	21.1	1.31 (0.80–2.14)	17	22.4	1.23 (0.77–1.97)	0.421
Drinking water source																	
Piped	397	84.6	21	5.3	1	1.000	9	2.3	1	1.000	65	16.4	1	76	19.1	1	
Rudimentary well	36	7.7	4	11.1	1.99 (0.72–5.99)	0.259	2	5.6	2.32 (0.52–10.35)	0.248	9	25.0	1.44 (0.78–2.67)	4	11.1	0.55 (0.22–1.42)	0.271
Water truck	17	3.6	1	5.9	1.11 (0.16–7.79)	0.612	–	–	Undefined	1.000	1	5.9	0.36 (0.05–2.43)	1	5.9	0.30 (0.04–2.07)	0.217
Direct from river	19	4.1	6	31.6	5.97 (2.73–13.0)	0.001	–	–	Undefined	1.000	4	21.1	1.28 (0.52–3.15)	7	36.8	1.92 (1.03–3.58)	0.075

Conventional sign used: – Data equal to zero, not resulting from rounding; PR: Prevalence ratio; 95% CI: Confidence intervals.



Dotted lines delimit heavy ($\geq 4,000$ epg), moderate (2,000 to 3,999 epg), and light ($\leq 1,999$ epg) parasitic loads.

Figure 2 – Scatter diagram demonstrating the parasite loads of hookworm infections in relation to age and sex of infected residents in Canto Grande and Alto Bonito communities, Chapada das Mesas National Park, Maranhão State, Brazil, 2022

Of 59 hookworm-positive subjects, 50 had their infection intensity assessed by Kato-Katz smears. The parasitic loads detected ranged from 24 epg to 16,728 epg, with an average of 1,304 epg (standard deviation = 417 epg). It was observed that 86.0% ($n = 43$) of hookworm-positive subjects had light, 4.0% ($n = 2$) had moderate, and 10.0% ($n = 5$) had heavy hookworm infection intensity (Figure 2). From seven *A. lumbricoides* infections identified, the parasitic load of three could be determined with two moderate and one light infection detected.

Figure 3 shows the spatial distribution of intestinal parasites in the studied communities. It was observed that the STH hotspots were in the Canto Grande (Carolina) community, while the protozoa hotspots were concentrated in Alto Bonito (Riachão).

DISCUSSION


The present study assessed the epidemiology of intestinal parasitism in rural communities living around an environmental protection area of Maranhão State, northeastern Brazil. As assessed by the socio-ecological data survey, almost half of the population lives in poverty and almost 1/3 practices open defecation, which characterizes a favorable scenario to the spread of parasitic diseases. The main finding is the persistence of STH transmission foci, mainly hookworms, significantly associated with open defecation in the peri-domestic environment. Therefore, differences in the vulnerability to intestinal parasitism could be observed

on the micro-regional scale. Studies in traditional communities carried out in other countries have also demonstrated the vulnerability of these populations to intestinal parasitic infections and increased risk in demographic concentration and poverty situations^{19,20}. The prevalence of hookworm infection was significantly higher in the Canto Grande community, which is predominantly rural and, as observed in fieldwork, has poorer sanitation conditions. On the other hand, ascariasis and giardiasis were significantly more frequent in Alto Bonito, which points to different transmission dynamics for parasites transmitted percutaneously and orally.

The transmission of hookworms is intrinsically related to soil contamination in peri-domestic environments with fecal matter. Previous studies conducted in the communities of Chapada das Mesas National Park demonstrated that most people practice open defecation⁶. It can be inferred that low sanitation, associated with poverty, allows the maintenance of hookworm transmission in the region. The association between open defecation and STH has been demonstrated in several regions, including Ethiopia^{21,22}, India²³, Mozambique²⁴, and Central America²⁵. Effective strategies to reduce the proportion of the population that practices open defecation are a major challenge in the region, where people maintain ancestral habits regarding how they live and interact with the environment. In this sense, the study data, which have a micro-regional scope, can be extrapolated to different communities in similar socio-ecological contexts and where open defecation

is still practiced. Solutions to interrupt the transmission of intestinal parasites in traditional rural communities require more comprehensive approaches, focusing on health education, awareness of the relationship with the environment, and more efficient intervention strategies in the context of primary health care. The development of latrines adapted to rural communities in northeastern Brazil to improve sanitation and reduce environmental contamination with fecal matter remains a difficult task^{26,27,28,29,30}.

In the present study, although most of the STH infections were of light intensity, individuals with moderate and heavy parasitic loads of hookworms were identified, which demonstrates that these nematodes are a health problem in these communities and a potential cause of iron-deficiency anemia. It is important to note that the single-dose albendazole mass treatment strategy implemented in Brazil targets school-aged children, so infections in adults and preschoolers may feed the cycles of STH transmission in the studied communities.



The factors associated with hookworm infection were not related to infections by intestinal protozoa. Thus, a more uniform distribution was observed for these organisms, and the water consumption directly from a river was significantly associated with giardiasis. Previous studies in these communities have shown that the majority of residents of the National Park drink water collected directly from rivers, which are vulnerable to pollution from garbage and solid waste, in a scenario conducive to the transmission of waterborne diseases. Improving the management of drinking water supplies and food hygiene can influence intestinal protozoa transmission in the region. These organisms, which are not the target of any specific policy, have not been detected due to the unreliability of the most common diagnostic techniques and are more difficult to treat since single-dose drugs, such as secnidazole, are not yet universally available. Chronic giardiasis, apparently asymptomatic and thus often undiagnosed, impacts the physical development of children given its association with protein-caloric malnutrition^{31,32}. Data from this study reinforces that intestinal parasitism should be considered a heterogeneous group of infections and that the simplification of the phenomenon, which tends to contribute to the development of unified control strategies, should be avoided.

CONCLUSION

The present study concludes that STH and intestinal protozoa persist in traditional communities in Brazil, where improving sanitation systems is a challenge. Traditional ways of living and interacting with

the environment can be associated with the transmission of parasitic diseases perpetuation in a poverty scenario, lack of health education, absence of health promotion programs, and inefficient primary health care systems.

ACKNOWLEDGMENTS

We thank Carolina and Riachão administrative staff and Primary Health Care Agents for their support in field activities. We would also like to acknowledge the ICMBio for the logistical support provided during the home visits when planning the study.

FINANCIAL SUPPORT

This study was supported by the Laboratory of Innovations in Therapies, Education, and Bioproducts, the Oswaldo Cruz Institute, Fiocruz, Rio de Janeiro, and the Regional Office Fiocruz Piauí.

AUTHORS' CONTRIBUTIONS

MLAS, FACC, and AHAMN designed the study, analyzed the data, and wrote the manuscript. KJLM supervised the laboratory and field activities. CMOGA, JS, and MMA carried out fieldwork, including visits to homes and laboratory analyses. JCPB critically reviewed the data analysis and manuscript. The final manuscript was reviewed and approved by all authors.

CONFLICTS OF INTEREST

The authors declared no potential conflicts of interest concerning the research, authorship, and/or publication of this article.



REFERENCES

- 1 Brasil. Decreto s/n, de 12 de dezembro de 2005. Cria o Parque Nacional da Chapada das Mesas, nos municípios de Carolina, Riachão e Estreito, no estado do Maranhão, e dá outras providências. Diário Oficial da União [Internet]. Brasília (DF), 2005 dez 13 [citado 2022 jul 21]; Seção 1:2. Disponível em: <https://pesquisa.in.gov.br/imprensa/jsp/visualiza/index.jsp?jornal=1&pagina=2&data=13/12/2005>.
- 2 Amarante CB. Reconhecimento jurídico-normativo das populações tradicionais pelo estado brasileiro: uma revisão. Encicl Biosfera. 2011;7(12):1-9.
- 3 Arruda R. "Populações tradicionais" e a proteção dos recursos naturais em unidades de conservação. Ambient Soc. 1999 dez;5:79-92.
- 4 Silva MLA, Paiva LS, Araújo MFV, Conceição GM. Environmental perception of residents of the Parque Nacional da Chapada das Mesas, in the phytogeographic domain in the forests from Brazilian Cerrado. Rev Espacios. 2017;38(22):33-40.
- 5 Silva MLA, Araújo MFV, Conceição GM. Chapada das Mesas National Park (Maranhão/Brazil): socio-economic activities of dwellers and their reflexes. Rev Ibero-Am Cienc Ambient. 2020 Fev-Mar;11(2):381-92.
- 6 Silva MLA, Araújo MFV, Conceição GM. Identity and way of life of the dwellers of the region of Chapada das Mesas, Maranhão, Brazil. Biota Amazon. 2017;7(4):41-7.
- 7 Haque R. Human intestinal parasites. J Health Popul Nutr. 2007 Dec;25(4):387-91.
- 8 Jourdan PM, Lamberton PHL, Fenwick A, Addiss DG. Soil-transmitted helminth infections. Lancet. 2018 Jan;391(10117):252-65.
- 9 Vivancos V, González-Alvarez I, Bermejo M, Gonzalez-Alvarez M. Giardiasis: Characteristics, pathogenesis and new insights about treatment. Curr Top Med Chem. 2018;18(15):1287-303.
- 10 Carrero JC, Reyes-López M, Serrano-Luna J, Shibayama M, Unzueta J, León-Sicairos N, et al. Intestinal amoebiasis: 160 years of its first detection and still remains as a health problem in developing countries. Int J Med Microbiol. 2020 Jan;310(1):151358.

- 11 Sard BG, Navarro RT, Esteban Sanchis JG. Non-pathogenic intestinal amoebae: a clinical-analytical overview. *Enferm Infecc Microbiol Clin*. 2011 Mar;29(Suppl 3):20-8.
- 12 Echazú A, Bonanno D, Juarez M, Cajal SP, Heredia V, Caropresi S, et al. Effect of poor access to water and sanitation as risk factors for soil-transmitted helminth infection: selectiveness by the infective route. *PLoS Negl Trop Dis*. 2015 Sep;9(9):e0004111.
- 13 Ferreira FS, Baptista-Fernandes T, Oliveira D, Rodrigues R, Neves E, Lima A, et al. *Giardia duodenalis* and soil-transmitted helminths infections in children in São Tomé and Príncipe: do we think *Giardia* when addressing parasite control? *J Trop Pediatr*. 2015 Apr;61(2):106-12.
- 14 Calegar DA, Nunes BC, Monteiro KJL, Bacelar PAA, Evangelista BBC, Almeida MM, et al. Genotypic and epidemiologic profiles of *Giardia duodenalis* in four Brazilian biogeographic regions. *Microorganisms*. 2022 Apr;10(5):940.
- 15 Katz N. Inquérito nacional de prevalência da esquistossomose mansoni e geo-helmintoses. Belo Horizonte: CPqRR; 2018. (Série Esquistossomose; 17).
- 16 Almeida MM, Monteiro KJL, Bacelar PAA, Santos JP, Freitas SPC, Evangelista BBC, et al. Interactions between malnutrition, soil-transmitted helminthiasis and poverty among children living in periurban communities in Maranhão State, Northeastern Brazil. *Rev Inst Med Trop S Paulo*. 2020;62:e73.
- 17 QGIS Development Team. QGIS Geographic Information System. Open Source Geospatial Foundation Project [Internet]. 2023 [cited 2021 Apr 27]. Available from: <http://qgis.osgeo.org>.
- 18 Pan American Health Organization, World Health Organization. Operational guidelines for the implementation of integrated deworming activities: a contribution to the control of soil-transmitted helminth infections in Latin America and the Caribbean. Washington: PAHO, WHO; 2015.
- 19 Muslim A, Mohd Sofian S, Shaari SA, Hoh BP, Lim YA. Prevalence, intensity and associated risk factors of soil transmitted helminth infections: a comparison between Negritos (indigenous) in inland jungle and those in resettlement at town peripheries. *PLoS Negl Trop Dis*. 2019 Apr;13(4):e0007331.
- 20 Anegagrie M, Lanfri S, Aramendia AA, Scavuzzo CM, Herrador Z, Benito A, et al. Environmental characteristics around the household and their association with hookworm infection in rural communities from Bahir Dar, Amhara Region, Ethiopia. *PLoS Negl Trop Dis*. 2021 Jun;15(6):e0009466.
- 21 Muluneh C, Hailu T, Alemu G. Prevalence and associated factors of soil-transmitted helminth infections among children living with and without open defecation practices in northwest Ethiopia: a comparative cross-sectional study. *Am J Trop Med Hyg*. 2020 Jul;103(1):266-72.
- 22 Aemiro A, Menkir S, Tegen D, Tola G. Prevalence of soil-transmitted helminthes and associated risk factors among people of Ethiopia: a systematic review and meta-analysis. *Infect Dis (Auckl)*. 2022 Mar;15:1-20.
- 23 Greenland K, Dixon R, Khan SA, Gunawardena K, Kihara JH, Smith JL, et al. The epidemiology of soil-transmitted helminths in Bihar State, India. *PLoS Negl Trop Dis*. 2015 May;9(5):e0003790.
- 24 Casmo V, Augusto G, Nala R, Sabonete A, Carvalho-Costa FA. The effect of hookworm infection and urinary schistosomiasis on blood hemoglobin concentration of schoolchildren living in northern Mozambique. *Rev Inst Med Trop S Paulo*. 2014 May-Jun;56(3):219-24.
- 25 Krause RJ, Koski KG, Pons E, Sandoval N, Sinisterra O, Scott ME. *Ascaris* and hookworm transmission in preschool children from rural Panama: role of yard environment, soil eggs/larvae and hygiene and play behaviours. *Parasitology*. 2015 Oct;142(12):1543-54.
- 26 Calegar DA, Bacelar PAA, Evangelista BBC, Monteiro KJL, Dos Santos JP, Almeida MM, et al. Socioenvironmental factors influencing distribution and intensity of soil-transmitted helminthiasis in the Brazilian Amazon: Challenges for the 2030 Agenda. *J Trop Med*. 2021 Feb;2021:6610181.
- 27 Kligerman DC, Cardoso TAO, Cohen SC, Azevedo DCB, Toledo GA, Azevedo APCB, et al. Methodology for a comprehensive health impact assessment in water supply and sanitation programmes for Brazil. *Int J Environ Res Public Health*. 2022 Oct;19(19):12776.
- 28 Barreto ML, Genser B, Strina A, Teixeira MG, Assis AMO, Rego RF, et al. Impact of a citywide sanitation program in Northeast Brazil on intestinal parasites infection in young children. *Environ Health Perspect*. 2010 Nov;118(11):1637-42.
- 29 Sousa ACA, Costa NR. Basic sanitation policy in Brazil: discussion of a path. *Hist Cienc Saude-Manguinhos*. 2016 Jul-Sep;23(3):615-34.
- 30 Raupp L, Cunha GM, Fávoro TR, Santos RV. Sanitation conditions of indigenous and nonindigenous households in Brazil according the 2000 and 2010 national censuses. *Cienc Saude Coletiva*. 2020 Oct;25(10):3753-63.

- 31 Coronato-Nunes B, Calegar DA, Monteiro KJL, Hubert-Jaeger L, Reis ERC, Xavier SCC, et al. *Giardia intestinalis* infection associated with malnutrition in children living in northeastern Brazil. *J Infect Dev Ctries*. 2017 Jul;11(7):563-70.
- 32 Carvalho-Costa FA, Gonçalves AQ, Lassance SL, Silva Neto LM, Salmazo CAA, Bóia MN. *Giardia lamblia* and other intestinal parasitic infections and their relationships with nutritional status in children in Brazilian Amazon. *Rev Inst Med Trop S Paulo*. 2007 May-Jun;49(3):147-53.

Received / Recebido em: 27/6/2023

Accepted / Aceito em: 19/4/2024

Este artigo passou por atualização. Ver **Errata** publicada em: <https://doi.org/10.5123/S2176-6223202401645>.

How to cite this article / Como citar este artigo:

Silva MLA, Álvares CMOG, Silva J, Bordignon JCP, Almeida MM, Carneiro ELNC, et al. Intestinal parasitism in traditional communities surrounding an environmental protection area in Maranhão State, northeastern Brazil. *Rev Pan Amaz Saude*. 2024;15:e202401484. Doi: <https://doi.org/10.5123/S2176-6223202401484>